

TABLE 1

Analyte	Maplewood Acceptance (revised 6/29//2015)	Farms Criteria
GC/MS SEMI VOC/PAHs BY 8270D (mg/kg)		
2-Methylnaphthalene	<0.7	
Acenaphthene	<4	
Acenaphthylene	<1	
Anthracene	<10	
Benzo[a]anthracene	<7	
Benzo[a]pyrene	<2	
Benzo[b]fluoranthene	<7	
Benzo[g,h,i]perylene	<10	
Benzo[k]fluoranthene	<10	
Chrysene	<20	
Dibenz(a,h)anthracene	<0.7	
Fluoranthene	<40	
Fluorene	<10	
Indeno[1,2,3-cd]pyrene	<7	
Naphthalene	<4	
Phenanthrene	<10	
Pyrene	<40	
GC/MS VOA BY 8260C (mg/kg)		
ALL VOCs	not detectable	
GC SEMI VOA BY 8100 Modified		
TPH (C9-C36)	<500	
PCBs BY 8082A (mg/kg)		
PCBs	<0.1	
Pesticides BY 8081B (mg/kg)		
	ND	
Herbicides BY 8151A (mg/kg)		
	ND	
METALS BY 6010C (mg/kg)		
Arsenic	<20	
Barium	<375	
Cadmium	<20	
Chromium (III)	<225	
Chromium (VI)	<100	
Chromium (Total)	<225	
Lead	<200	
Mercury	<3	
Selenium	<5	
Silver	<6	

TABLE 1

Analyte	Maplewood Acceptance (revised 6/29/2015)	Farms Criteria
OTHER METALS (mg/kg)		
ALUMINUM	<25000	
ANTIMONY	<10	
BERYLLIUM	<4	
COBALT	<40	
COPPER	<300	
IRON	<50000	
MAGNESIUM	<12500	
MANGANESE	<1500	
NICKEL	<150	
THALLIUM	<6	
VANADIUM	<225	
ZINC	<500	
TOXICITY CHARACTERISTIC LEACHING PROCEDURE LEAD (mg/l)		
TCLP LEAD	<5	
GENERAL CHEMISTRY BY SM 2510B (umhos/cm)		
Specific Conductance	<2000	
pH (S.U.)	5.0-9	
REACTIVE CN (mg/kg)	non reactive	
REACTIVE SULFIDE (mg/kg)	non reactive	
FLASHPOINT (degree F)	absent	
IGNITABILITY (degree F)	absent	

Samples should be collected at a frequency of one per 500 cy

Samples should be screened with a PID. Anything greater than 2 ppmV is not acceptable